

Prevention is better than the cure, but the cure cannot be worse than the disease: fiberoptic tracheal intubation in COVID-19 patients

Massimiliano Sorbello^{1,*}, Ida Di Giacinto², Ruggero M. Corso³, Rita Cataldo⁴, on behalf of the Società Italiana di Anestesia Analgesia Rianimazione e Terapia Intensiva (SIAARTI) Airway Management Research Group

¹Catania, Italy, ²Bologna, Italy, ³Forlì, Italy and ⁴Rome, Italy

*Corresponding author. E-mail: maxsorbello@gmail.com

Keywords: aerosol-generating procedure; airway management; COVID-19; fiberoptic tracheal intubation; high-flow nasal oxygen; rapid sequence induction

Editor—We would like to comment on the paper by Wu and colleagues¹ in light of the protocols and clinical recommendations² produced during the coronavirus disease 2019 (COVID-19) outbreak in Italy. COVID-19-related hypoxaemia and desaturation patterns are atypical, and might not be comparable with typical respiratory failure or acute respiratory distress syndrome (ARDS). Patients exhibit a low P_{aO_2}/F_{iO_2} (P/F) ratio with (initially) preserved high pulmonary compliance and shunt fractions, the so-called silent hypoxia,³ which has implications for airway management. In this respect, we note the following concerns with the study by Wu and colleagues¹:

- (i) The power analysis was calculated using 23% desaturation estimated on the basis of ICU patients, whereas COVID-19 patients desaturate further and faster. In their study, with initial mean (inter-quartile range [IQR]) P/F ratios of 139.5 (118.3–162.3) and 128.5 (121.5–136.3), the authors report mean saturations of 94% and 91%, respectively, with a range of 86–95.8%.¹ In daily practice, with an intubation trigger around P/F=150 mm Hg, COVID-19 patients quickly reach much lower Sp_{O_2} values upon apnoea.²
- (ii) ‘Rapid-sequence fiberoptic bronchoscopic tracheal intubation’¹ appears an oxymoron: fiberoptic intubation requires skill and is time consuming (72–420 s time range for awake intubation⁴). Wu and colleagues¹ report mean (IQR) values of 68.5 s (62.2–74.0 s) and 76.0 (68.0–90.5 s) for fiberoptic intubation in anaesthetised patients, probably reflecting the skill of the intubating team and the impact of anaesthesia. For a fast procedure, it is still too long not to result in significant desaturation, particularly considering that no oxygen was administered in the control group (in an obvious study comparing oxygen vs non-oxygen), and that, ‘There was no significant difference in the proportion of patients with minimum $Sp_{O_2} >95\%$ during intubation, in the incidence of $Sp_{O_2} <80\%$ during intubation’.¹ In other words, this might mean that, in a 30+30 COVID-19 patient sample, either extra oxygen is not useful or that high-flow nasal oxygen (HFNO) does not work.
- (iii) We do not believe that ‘HFNO shortened the duration of intubation. [...] but one possibility is that interruption of attempts at tracheal intubation to carry out rescue face-

mask ventilation was less frequently required in the HFNO group’. High-flow nasal oxygen probably reduced the number of attempts at rescue face-mask ventilation, and not the time for fiberoptic intubation, keeping in mind that the control group was not given oxygen.

Fiberoptic intubation is considered the gold standard for predicted difficult intubation, the underlying concept being that maintenance of spontaneous breathing preserves oxygenation.⁵ Nevertheless, use of a technique requiring longer times for intubation, with higher likelihood of desaturation and need for rescue face-mask ventilation (1+8 cases in the series of Wu and colleagues¹) cannot be recommended in COVID-19 patients made apnoeic by neuromuscular blocking agents.

- (iv) The authors report that, ‘During attempts at tracheal intubation, HFNO was maintained for the HFNO group’,¹ but they did not clarify if intubation was performed by the nasal or oral route. In the first case, either the 4.5 mm fibroscope was inserted in the nostril together with the HFNO prong (thus, resulting in a difficult scope movement), or in place of the HFNO prongs. Fiberoptic intubation is known to be more challenging via the oral route in anaesthetised patients because of airway collapse, especially if a dedicated airway is not used.⁶ If the nasal route was chosen, removing the HFNO prongs to allow passage of the bronchoscope would have reduced the benefit of apnoeic HFNO and increased aerosolisation risk.^{2,7–9}
- (v) Despite being superior in preventing hypoxaemia during rapid sequence induction and intubation, HFNO is not recommended in COVID-19 patients^{2,8,9} as (awake) fiberoptic intubation is considered an aerosol-generating procedure. It is then paradoxical to state that, ‘Rapid-sequence fiberoptic bronchoscopic tracheal intubation in patients with COVID-19 pneumonia may reduce the risk of viral spread’,¹ especially if the declared advantage is, ‘an increase of the distance between anaesthesiologist and patient’s airway’.¹ Bronchoscopes with an average length of 65 cm do not allow the 1 m distance considered safe for airborne spreading.^{2,7–9}

A videolaryngoscope (especially with external screen) works with similar distance, but has been shown to be faster and to require lower skills⁴ and to have simpler options for

rescue intervention.^{1,2} The same message comes from Chinese experience⁶ and from worldwide recommendations.^{2,8,9} To maximise first-pass success, we recommend a preloaded bougie or stylet, rapid sequence induction with full-dose neuromuscular blockade, preoxygenation by continuation of ongoing noninvasive ventilation, and apnoeic low-flow (1–3 L min⁻¹) oxygen through standard nasal prongs (nasal oxygen during efforts securing a tube [NODESAT]).² After two failed laryngoscopies, we recommend rescue use of fiberoptic intubation only through a second-generation intubating supraglottic airway device, which allows ventilation with limited environmental contamination.²

Airway management is complex in COVID-19 patients (infection risk, use of personal protective equipment (PPE), difficult communication, rapidly deteriorating patients, and shunt–hypoxaemia). Given the duration, complexity, and aerosolisation potential of fiberoptic intubation and the potential low-efficacy/high contamination profile of HFNO, we strongly discourage the use of the technique proposed by Wu and colleagues¹ in paralysed COVID-19 patients. Awake fiberoptic intubation remains the gold standard for predicted intubation/ventilation difficulty to be used in very selected cases in COVID-19 patients.^{2,8,9} Oxygenation, independently of disease, remains the main target of any airway management strategy,⁵ and although difficult at times, science and good sense should always prevail.

Declarations of interest

MS has received paid consultancy from Teleflex Medical, Verathon Medical, and DEAS Italia; is a patent co-owner (no royalties; DEAS Italia); and has received lecture grants and travel reimbursements (MSD Italia and MSD USA). IDG has received lecture grants and travel reimbursements from MSD Italia. There are no other competing interests declared.

References

1. Wu CN, Xia LZ, Li KH, et al. High-flow nasal-oxygenation-assisted fiberoptic tracheal intubation in critically ill patients with COVID-19 pneumonia: a prospective randomised controlled trial. *Br J Anaesth* 2020 Mar 19. <https://doi.org/10.1016/j.bja.2020.02.020>. pii: S0007-0912(20)30135-5 [Epub ahead of print].
2. Sorbello M, El-Boghdady K, Di Giacinto I, et al. The Italian coronavirus disease 2019 outbreak: experiences and recommendations from clinical practice. *Anaesthesia* 2020. <https://doi.org/10.1111/anae.15049>. Advance Access published on March 27
3. Ottestad W, Søvik S. COVID-19 patients with respiratory failure: what can we learn from aviation medicine? *Br J Anaesth* april 12, 2020. <https://doi.org/10.1016/j.bja.2020.04.012> [Epub ahead of print].
4. Jiang J, Ma DX, Li B, Wu AS, Xue FS. Videolaryngoscopy versus fiberoptic bronchoscope for awake intubation—a systematic review and meta-analysis of randomized controlled trials. *Ther Clin Risk Manag* 2018; **14**: 1955–63
5. Sorbello M, Afshari A, De Hert S. Device or target? A paradigm shift in airway management: implications for guidelines, clinical practice and teaching. *Eur J Anaesthesiol* 2018; **35**: 811–4
6. Meng L, Qiu H, Wan L, et al. Intubation and ventilation amid the COVID-19 outbreak: Wuhan's experience. *Anesthesiology* 2020. <https://doi.org/10.1097/ALN.0000000000003296>
7. Tran DT, Newton EK, Mount VA, Lee JS, Wells GA, Perry JJ. Rocuronium versus succinylcholine for rapid sequence induction intubation. *Cochrane Database Syst Rev* 2015; **10**: CD002788
8. Wax RS, Christian MD. Practical recommendations for critical care and anesthesiology teams caring for novel coronavirus (2019-nCoV) patients. *Can J Anaesth* 2020; **67**: 568–76
9. Peng PWH, Ho P-L, Hota SS. Outbreak of a new coronavirus: what anaesthetists should know. *Br J Anaesth* 2020; **124**: 497–501

doi: 10.1016/j.bja.2020.04.077

Advance Access Publication Date: 27 April 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

“Water, water, everywhere”: a challenge to ventilators in the COVID-19 pandemic

James Fullick^{1,*} and Michael Oliver²

¹Newport, UK and ²Cardiff, UK

*Corresponding author. E-mail: james.fullick@nhs.net

Keywords: COVID-19; pandemic; resource management; ventilator; water trap

Editor—With the spread of coronavirus disease 2019 (COVID-19), intensive care facilities have been rapidly overwhelmed across the UK and elsewhere.¹ In general, the UK

has fewer doctors and fewer ICU beds per capita than most of Europe.² Many hospitals have spread into the recovery unit of theatres and are using anaesthetic machines to