

CORRIGENDUM

Corrigendum to “Spinal anaesthesia for patients with coronavirus disease 2019 and possible transmission rates in anaesthetists: retrospective, single centre, observational cohort study” (Br J Anaesth 2020;124: 670-5)

Qi Zhong, Yin Y. Liu, Qiong Luo, Yu F. Zou, Hai X. Jiang, Hui Li, Jing J. Zhang, Zhen Li, Xin Yang, Min Ma, Li J. Tang, Ying Y. Chen*, Feng Zheng*, Jian J. Ke* and Zong Z. Zhang*

Department of Anaesthesiology, Zhongnan Hospital, Wuhan University, Wuhan, Hubei, China

*Corresponding authors. E-mails: chenyinyg@whu.edu.cn, fengzheng@whu.edu.cn, 1219628972@qq.com, zhangzz@whu.edu.cn
DOI of original article: <https://doi.org/10.1016/j.bja.2020.03.007>.

The authors regret that an error was present in Table 1. The corrected table is shown below:

The authors would like to apologise for any inconvenience caused.

Table 1 Baseline characteristics of 49 surgery patients with SARS-CoV-2 infection.

Characteristics	Number
Age, median (IQR), yr	31 (29–34)
Female, sex (%)	45 (91.8)
BMI, median (IQR), kg m ⁻²	35.2 (33.25–36.4)
Duration from onset of symptoms to radiological confirmation of pneumonia, median (IQR), days	3 (5–8)
Duration from onset of symptoms to operation room, median (IQR), days	2 (0.5–3)
Symptom (%)	
Cough	21 (42.8)
Sore throat	14 (28.6)
Myalgia	6 (12.2)
Shortness of breath	4 (8.2)
Gastrointestinal	2 (4.0)
Fever (%)	
<37.3°C	23 (46.9)
37.3°C–37.9°C	21 (42.9)
38.0°C–38.9 °C	3 (6.1)
>=39°C	2 (4.1)
Positive RT-PCR for COVID-19	13 (26.5)
Comorbidities (%)	
Hypertension	4 (8.2)
Diabetes mellitus	2 (4.1)
Thyroid dysfunction	2 (4.1)
Operation type (%)	
Caesarean section	45 (91.8)
Orthopedic operation	4 (8.2)

Continued

Table 1 Continued

Characteristics	Number
ASA physical status (%)	
1	43 (87.8)
2	6 (12.2)
Treatment (%)	
Nasal cannula	49 (100)
Antiviral agents	29 (59.2)
Antibacterial agents	47 (95.9)
Progress to severe pneumonia (%)	
Yes	0 (0)
No	100 (100)
Complication (%)	
Vomiting	3 (6.1)