

worsening respiratory conditions and leading to urgent ECMO' does not sound correct. Lack of preparation, use of a direct laryngoscope as a primary intubation device, and inappropriate bougie use likely all contributed to such an outcome. This simple device remains a precious adjunct for airway management when carefully used in conjunction with direct or videolaryngoscopy.¹⁰

Declarations of interest

MS has received paid consultancy from Teleflex Medical, Verathon Medical, and DEAS Italia; is a patent co-owner (no royalties) of DEAS Italia; and has received lecture grants and travel reimbursements from MSD Italia and MSD USA. GF has received royalties from Cook Medical (Frova Introducer; Bloomington, IN, USA) and Teleflex Medical (PercuTwist and EasyCric) for his airway device inventions. IH and GC declare no competing interests.

References

1. Abou-Arab O, Huette P, Berna P, Mahjoub Y. Tracheal trauma after difficult airway management in morbidly obese patients with COVID-19. *Br J Anaesth* 2020; **125**: e168–70
2. Petrini F, Di Giacinto I, Cataldo R, et al. Perioperative and periprocedural airway management and respiratory safety for the obese patient: 2016 SIAARTI Consensus. *Minerva Anestesiol* 2016; **82**: 1314–35
3. Sorbello M, El-Boghdady K, Di Giacinto I, et al. The Italian coronavirus disease 2019 outbreak: recommendations from clinical practice. *Anaesthesia* 2020; **75**: 724–32
4. Memtsoudis SG, Ivascu NS, Pryor KO, Goldstein PA. Obesity as a risk factor for poor outcome in COVID-19 induced lung injury: the potential role of undiagnosed obstructive sleep apnoea. *Br J Anaesth* 2020; **125**: e262–3
5. Sorbello M, Frova G. Frova introducer: neither a stylet nor simply an introducer. *Anaesthesia* 2008; **63**: 1010–1
6. Hodzovic I, Latto IP, Wilkes AR, Hall JE, Mapleson WW. Evaluation of Frova, single-use intubation introducer, in a manikin. Comparison with Eschmann multiple-use introducer and Portex single-use introducer. *Anaesthesia* 2004; **59**: 811–6
7. Yao W, Wang T, Jiang B, et al. Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *Br J Anaesth* 2020; **125**: e28–37
8. Frova G, Sorbello M. Iatrogenic tracheobronchial ruptures: the debate continues. *Minerva Anestesiol* 2011; **77**: 1130–3
9. Paul A, Gibson AA, Robinson OD, Koch J. The traffic light bougie: a study of a novel safety modification. *Anaesthesia* 2014; **69**: 214–8
10. Driver BE, Prekker ME, Klein LR, et al. Effect of use of a bougie vs endotracheal tube and stylet on first-attempt intubation success among patients with difficult airways undergoing emergency intubation: a randomized clinical trial. *JAMA* 2018; **319**: 2179–89

doi: 10.1016/j.bja.2020.05.013

Advance Access Publication Date: 19 May 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Airway management lessons from case reports of negative outcomes. Comment on *Br J Anaesth*; 125: e168–70

Thomas Heidegger

Grabs, Switzerland

E-mail: thomas.heidegger@srrws.ch

Keywords: airway management; case report; COVID-19; fiberoptic intubation; obesity; tracheal intubation

Always note and record the unusual ... Publish it. Place it on permanent record as a short, concise note. Such communications are always of value (Sir William Osler).¹

Editor—A forthcoming issue of the *British Journal of Anaesthesia* (BJA) contains a report of two cases of tracheal trauma after difficult airway management in morbidly obese patients with coronavirus disease 2019 (COVID-19).² The authors should be commended for submitting these negative outcomes to the BJA, and the BJA must be congratulated for publishing them. High-ranking journals rarely accept such

reports nowadays, and even though these reports often have major limitations, such as possible over-interpretation, lack of generalisability, or retrospective design,³ we can still learn a lot from them.⁴

Here are six lessons from that report:

- (i) *Airway management is (still) not as safe as we might believe.* It is rare that medical professionals report their own adverse events for at least two reasons: fear of reputational consequences and fear of legal consequences. Thus, there is likely a strong publication bias favouring case reports with positive results. Regarding safety of airway management,

results are likely too optimistic as well.⁴ A good example is the interpretation of the results from The Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society (NAP4), the largest prospective study of major airway events occurring during anaesthesia, in the ICU and emergency department.⁵ The authors themselves state that only roughly 25% of the cases with adverse events were captured, indicating a more accurate incidence of perhaps 1:5500 or even greater instead of the published incidence of 1:22 000 for major airway events.⁶ Identifying the true incidence of adverse airway events is theoretically simple: we need further large prospective data including all cases, both elective and emergency, and cases with good and adverse outcomes, including death.⁷

- (ii) *Obesity itself is a predictor of difficult airway management.* This was shown by the results of NAP4, in which obese patients were twice as common in the population that suffered incidents than in the group that did not,⁵ and this was confirmed in a recent study from Australia.⁸ Anatomical alterations, such as the combination of a large tongue and excessive upper airway soft tissue, and physiological alterations, such as decreased functional residual capacity and increased oxygen consumption, are important disadvantages for obese patients.⁹ Whether obesity is a predictor of difficult mask ventilation is still debated.¹⁰ Regarding tracheal intubation, an analysis of electronic records of more than 67 000 patients showed that a BMI of $>30 \text{ kg m}^{-2}$ was significantly associated with increased likelihood of more than one tracheal intubation attempt, but the odds of difficult intubation remained unchanged once BMI exceeded 30 kg m^{-2} .¹¹
- (iii) *Laryngoscopy is not the same as intubation.* It is important to distinguish between these two procedures, as difficult laryngoscopy can be followed by easy tracheal intubation. (This is common during conventional tracheal intubation.) Likewise, easy laryngoscopy can be followed by difficult tracheal intubation ('you see that you fail'), which is the most common cause of failed tracheal intubation with videolaryngoscopy.¹² Even though the use of videolaryngoscopy improves the glottic view, there is currently no evidence that it reduces the number of tracheal intubation attempts or the incidence of respiratory complications.¹³ Whether videolaryngoscopy would have prevented the adverse event in the cases described² remains unclear.
- (iv) *Tracheal tube introducers ('bougies') and rigid stylets are potentially dangerous.* The cases reported show the potential dangers of rigid tools, such as tracheal tube introducers or rigid stylets, to facilitate tracheal intubation in patients with COVID-19, particularly in the absence of glottis visibility.^{14,15}
- (v) *Fibreoptic intubation is an established alternative in patients with morbid obesity.* Even though fibreoptic intubation is infrequently used as a first-choice technique for management of patients with morbid obesity, it is a tried-and-tested alternative for such situations¹⁶ and should always be considered.
- (vi) *Airway management in obese patients with COVID-19: combining protection and best practice is essential.* Various national airway societies have published consensus guidelines addressing airway management during the COVID pandemic.^{17–19} Besides a few techniques that should be strictly avoided, such as high-flow nasal oxygen

and small-bore cannula cricothyroidotomy with jet ventilation, management of the difficult airway always involves weighing the risks for the patient and other persons involved, with the likelihood of first-attempt success. The technique chosen may ultimately differ according to local practices, resources, and experience.

In conclusion, publishing airway management case reports, above all those with negative outcomes, remains important. We need to expose more than the tip of the iceberg so that we can improve our daily practice in airway management to improve patient safety.

Declarations of interest

The author declares that they have no conflicts of interest.

Acknowledgements

The author thanks Jeannie Wurz, medical writer/editor, Bern, Switzerland.

References

1. Thayer WS. *Osler, the teacher*. Sir William Osler. Bart. Baltimore, MD: Johns Hopkins Press; 1920. 51–2
2. Abou-Arab O, Huette P, Berna P, Mahjoub Y. Tracheal trauma after difficult airway management in patients with COVID-19. *Br J Anaesth* 2020; **125**: e168–70
3. Nissen T, Wynn R. The clinical case report: a review of its merits and limitations. *BMC Res Notes* 2014; **7**: 264
4. Albrecht J, Meves A, Bigby M. Case reports and case series from *Lancet* had significant impact on medical literature. *J Clin Epidemiol* 2005; **58**: 1227–32
5. Cook TM, Woodall N, Frerk C. Major complications of airway management in the UK: results of the fourth national audit project of the royal college of anaesthetists and the difficult airway society. Part 1: anaesthesia. *Br J Anaesth* 2011; **106**: 617–31
6. O'Sullivan E, Laffey Y, Pandit JJ. A rude awakening after our fourth 'NAP': lessons for airway management. *Anaesthesia* 2011; **66**: 331–40
7. Devereaux PJ, Chan MT, Eisenach J, Schrickler T, Sessler DI. The need for large clinical studies in perioperative medicine. *Anesthesiology* 2012; **116**: 1169–75
8. Schnittker R, Marshall SD, Berecki-Gisolf J. Patient and surgery factors associated with the incidence of failed and difficult intubation. *Anaesthesia* 2020; **75**: 756–66
9. Murphy C, Wong DT. Airway management and oxygenation in obese patients. *Can J Anaesth* 2013; **60**: 929–45
10. Kheterpal S, Han R, Tremper KK, et al. Incidence and predictors of difficult and impossible mask ventilation. *Anesthesiology* 2006; **105**: 885–91
11. Saasouh W, Laffey K, Turan A, et al. Degree of obesity is not associated with more than one intubation attempt: a large centre experience. *Br J Anaesth* 2018; **120**: 1110–6
12. Aziz MF, Dillman D, Fu R, Brambrink AM. Comparative effectiveness of the C-MAC video laryngoscope versus direct laryngoscopy in the setting of the predicted difficult airway. *Anesthesiology* 2012; **116**: 629–36
13. Lewis SR, Butler AR, Parker J, Cook TM, Schofield-Robinson OJ, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal

- intubation: a Cochrane Systematic Review. *Br J Anaesth* 2017; **119**: 369–83
14. Hodzovic I, Latto IP, Wilkes AR, Hall JE, Mapleson WW. Evaluation of Frova, single-use intubation introducer, in a manikin. Comparison with Eschmann multiple-use introducer and Portex single-use introducer. *Anaesthesia* 2004; **59**: 811–6
 15. Sorbello M, Hodzovic, Cusumano G, Frova G. Tracheal introducers and airway trauma COVID-19. *Br J Anaesth* 2020; **125**: e168–70
 16. Heidegger T, Gerig HJ, Ulrich B, Schnider TW. Structure and process quality illustrated by fiberoptic intubation: analysis of 1612 cases. *Anaesthesia* 2003; **58**: 734–9
 17. Cook TM, El-Boghdadly K, McGuire B, McNarry AF, Patel A, Higgs A. Consensus guidelines for managing the airway in patients with COVID-19. *Anaesthesia* 2020; **75**: 785–99
 18. Patwa A, Shah A, Garg R, et al. All India Difficult Airway Association (AIDAA) consensus guidelines for airway management in the operating room during the COVID-19 pandemic. *Indian J Anaesth* 2020; **64**: S107–15
 19. Yao W, Wang T, Jiang B, et al. Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *Br J Anaesth* 2020; **125**: e28–37

doi: 10.1016/j.bja.2020.06.042

Advance Access Publication Date: 1 July 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Low-flow nasal cannula oxygen and potential nosocomial spread of COVID-19

Sara N. Goldhaber-Fiebert^{1,*}, Jeremy A. Greene² and Brian T. Garibaldi²

¹Stanford, CA, USA and ²Baltimore, MD, USA

*Corresponding author. E-mail: saragf@stanford.edu

Keywords: COVID-19; infection control; low-flow; nasal cannula; nosocomial spread; oxygen delivery; SARS-CoV-2

Editor—In the coronavirus disease 2019 (COVID-19) pandemic airway procedures such as intubation, noninvasive positive pressure ventilation, and high-flow nasal cannula are widely considered as potential risks for nosocomial transmission, and risks of infection are recognised even from asymptomatic patients.^{1–3} Yet to date there has been little published and limited awareness regarding the risks of a far more prevalent practice: low-flow nasal cannula oxygen spread of COVID-19 from unsuspected patients. Even detailed studies of clinician exposures to unsuspected COVID-19 patients frequently do not include low-flow nasal cannula oxygen therapy as an exposure category.⁴

High viral loads of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are present in human nares in symptomatic and asymptomatic patients.⁵ Air dispersion from low-flow nasal cannulae can reach more than 1 meter away, a distance which, although less than that of an uncovered cough, produces a constant rate of dispersal similar to that of noninvasive ventilation masks set at more than double the oxygen flows.⁶ Viral particles survive for multiple days on many surfaces, further facilitating nosocomial transmission.⁷ Even without aerosolisation, surface contamination risks remain. During the first SARS epidemic, supplemental oxygen therapy emerged as a risk factor for nosocomial transmission on open wards, equivalent to patient bed crowding and failure to provide washing stations for providers.⁸ Even with single-occupancy rooms, healthcare providers could be exposed to or spread SARS-CoV-2 after touching contaminated surfaces

surrounding unsuspected COVID-19 patients presenting for other reasons. In a recent study, researchers sampling air inside COVID-19 negative-pressure patient rooms found the highest concentration of viral RNA in the room of a patient who was on oxygen 1 L min⁻¹ by nasal cannula, with no documented cough,⁹ although in this context clinicians were wearing full protective equipment.

Some institutions have begun covering low-flow nasal cannulae, at least in certain contexts,^{10,11} although discussions with peers across specialities and institutions suggest that practice is far from uniform and is sometimes limited to known COVID-19 patients. Existing data should give institutions pause to consider the infection risks of oxygen delivery for all patients, especially in cases where oxygen use is informed by habit, rather than evidence of clinical benefit. When low-flow oxygen via nasal cannula is clearly indicated, simple strategies can be used to mitigate the risk of spread. For example before extubation, a nasal cannula can be placed and covered with a surgical mask to limit the potential for environmental contamination.¹¹

By a conservative estimate, if 10% of the occupants of the roughly 1 million hospital beds in the USA are on low-flow nasal cannula oxygen on any given day, that translates into 100,000 patients in US hospitals whose treatment may also be adding to nosocomial spread of SARS-CoV-2. Local conditions and supplies should guide considerations of using surgical masks to cover all low-flow nasal cannulae. If surgical masks are in short supply, other coverings,