

---

This article has been retracted: please see Elsevier Policy on Article Withdrawal (<https://www.elsevier.com/about/our-business/policies/article-withdrawal>).

This article has been retracted at the request of the Editor-in-Chief, Professor Hugh Hemmings, based on the recommendations of Justus-Liebig-University Giessen following an internal review of research conducted by Joachim Boldt at the University. This is further described in 'Further Retractions of Articles by Joachim Boldt', <https://doi.org/10.1016/j.bja.2020.02.024>.

*British Journal of Anaesthesia*, 125 (3): 413 (2020)

doi: [10.1016/j.bja.2020.07.029](https://doi.org/10.1016/j.bja.2020.07.029)

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

## Retraction notice to “The role of enoximone in cardiac surgery” [*Br J Anaesth* 69 (1992) 45–50]

J. Boldt<sup>1,\*</sup>, C. Knothe<sup>1</sup>, B. Zickmann<sup>1</sup>, M. Ballesteros<sup>1</sup>, W. Russ<sup>1</sup>, F. Dapper<sup>2</sup> and G. Hempelmann<sup>1</sup>

<sup>1</sup>Department of Anaesthesiology and Intensive Care Medicine, Justus-Liebig-University Giessen, D-6300 Giessen, Germany and <sup>2</sup>Clinic of Cardiovascular Surgery, Justus-Liebig-University Giessen, D-6300 Giessen, Germany

\*Corresponding author.

DOI of original article: <https://doi.org/10.1093/bja/69.1.45>.

---

This article has been retracted: please see Elsevier Policy on Article Withdrawal (<https://www.elsevier.com/about/our-business/policies/article-withdrawal>).

This article has been retracted at the request of the Editor-in-Chief, Professor Hugh Hemmings, based on the recommendations of Justus-Liebig-University Giessen following an internal review of research conducted by Joachim Boldt at the University. This is further described in 'Further Retractions of Articles by Joachim Boldt', <https://doi.org/10.1016/j.bja.2020.02.024>.

*British Journal of Anaesthesia*, 125 (3): 413–414 (2020)

doi: [10.1016/j.bja.2020.07.030](https://doi.org/10.1016/j.bja.2020.07.030)

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

## Retraction notice to “Volume replacement with hydroxyethyl starch solution in children” [*Br J Anaesth* 70 (1993) 661–5]

J. Boldt<sup>1,\*</sup>, C. Knothe<sup>1</sup>, E. Schindler<sup>1</sup>, H. Hammermann<sup>1</sup>, F. Dapper<sup>2</sup> and G. Hempelmann<sup>1</sup>

<sup>1</sup>Department of Anaesthesiology and Intensive Care Medicine, Justus-Liebig-University Giessen, D-6300 Giessen, Germany and <sup>2</sup>Department of Cardiovascular Surgery, Justus-Liebig-University Giessen, D-6300 Giessen, Germany

\*Corresponding author.

DOI of original article: <https://doi.org/10.1093/bja/70.6.661>.

---

This article has been retracted: please see Elsevier Policy on Article Withdrawal (<https://www.elsevier.com/about/our-business/policies/article-withdrawal>).