

Authors' contributions

Study design: HZ, BHS, LAC, JH
 Data collection: HZ, JT, XZ, AL, LW, WZ
 Writing of article: HZ, LAC
 Critical revision: XZ, HLH, WM, BHS, JH
 Approval of final version: all authors.

Declarations of interest

LAC is an editor of the *British Journal of Anaesthesia*. The other authors declare that they have no conflicts of interest.

Acknowledgements

University of Dundee Global Challenges Research Fund.

References

1. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; **382**: 727–33
2. WHO. Coronavirus disease 2019 (COVID-19) situation report-190 2020. Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200728-covid-19-sitrep-190.pdf?sfvrsn=fec17314_2. [Accessed 28 July 2020]
3. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020; **323**: 1239–42. <https://doi.org/10.1001/jama.2020.2648>
4. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020; **180**: 934–43. <https://doi.org/10.1001/jamainternmed.2020.0994>. [Accessed 13 March 2020]
5. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; **395**: 1054–62
6. Nepogodiev Dmitri, et al. Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet* 2020; **396**: 27–38
7. China's National Health Commission. *Diagnosis and treatment scheme of COVID-19 (Interim version 7)*. Available from: 2020. English version translated by the Chinese Society of Cardiology. Available from, <http://www.nhc.gov.cn/yzygj/s7653p/202003/46c9294a7dfe4cef80dc7f5912eb1989.shtml>. [Accessed 26 June 2020]. <http://kjfy.meetingchina.org/msite/news/show/cn/3337.html>
8. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; **382**: 1708–20
9. Dudley JP, Lee NT. Disparities in age-specific morbidity and mortality from SARS-CoV-2 in China and the Republic of Korea. *Clin Infect Dis* 2020; **71**: 863–5. <https://doi.org/10.1093/cid/ciaa354>
10. Mo X, Jian W, Su Z, et al. Abnormal pulmonary function in COVID-19 patients at time of hospital discharge. *Eur Respir J* 2020; **55**: 2001217. <https://doi.org/10.1183/13993003.01217-2020>. [Accessed 18 June 2020]
11. Moppett IK, Gardiner D, Harvey DJR. Guidance in an uncertain world. *Br J Anaesth* 2020; **125**: 7–9

doi: 10.1016/j.bja.2020.07.001

Advance Access Publication Date: 16 July 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Hazardous mismatch between pulmonary pathogens and antibiotic treatments in COVID-19 patients

Silvia Coppola¹, Arianna Ciabattoni², Tommaso Pozzi², Valentina Castagna², Gianluigi L. Bassi⁴ and Davide Chiumello^{1,2,3,*}

¹Department of Anesthesia and Intensive Care, ASST Santi Paolo e Carlo, San Paolo University Hospital, Milan, Italy, ²Department of Health Sciences, University of Milan, Milan, Italy, ³Coordinated Research Center on Respiratory Failure, University of Milan, Milan, Italy and ⁴Critical Care Research Group, The Prince Charles Hospital, Chermside, Australia

*Corresponding author. E-mail: davide.chiumello@unimi.it

Keywords: antibiotic therapy; bacterial pneumonia; coinfection; COVID-19; mechanical ventilation

Editor—In late December 2019, a new outbreak of pneumonia of unknown origin was described in Wuhan, China,¹ soon ascribed to the novel coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).² The associated coronavirus disease 2019 (COVID-19) results in respiratory failure in up to 90% of infected patients admitted to the ICU, and manifests with chest CT abnormalities such as ground glass opacities, and multifocal patchy consolidation with or without interstitial changes with a peripheral distribution.³ Most patients present additional organ dysfunction including acute respiratory distress syndrome (ARDS), septic shock, and cardiac, kidney and hepatic failure. Although there is no current cure for COVID-19, immunomodulatory agents, such as corticosteroids, tocilizumab (anti-interleukin-6 monoclonal antibody), and others have been tested to modulate the cytokine storm that often ensues from this novel viral infection.⁴ The main risk factor associated with these agents is the risk of superimposed pulmonary co-infections.

Given that the majority of COVID-19 patients present unspecific respiratory symptoms and chest radiography abnormalities, empirical antibiotic treatments are commonly prescribed before and during hospitalisation.⁵ The recent Surviving Sepsis Campaign guidelines on the management of critically ill patients with SARS-CoV-2 infection suggested empiric antimicrobials/antibacterial agents (as a weak recommendation with low-quality of evidence).⁶ Inflammatory markers, such as procalcitonin, have been applied for antimicrobial stewardship, yet in COVID-19, inflammatory biomarkers are generally elevated.

Only limited evidence has been published on pulmonary co-infections in COVID-19 patients undergoing mechanical ventilation. Hence, there is urgency to identify causative pathogens in this population to guide appropriate antibiotic therapy while reducing the risk of drug resistance. Herein, we report data from 53 COVID-19 patients admitted to the ICU and on mechanical ventilation in a single-centre study with a median age of 57 (9) yr. Among these, 26 (49%) received empiric antibiotic therapy upon hospital admission. As recommended for patients with community-acquired pneumonia,⁷ empirical therapy comprised ceftriaxone plus a macrolide or levofloxacin (in 17

patients, 32%), but also piperacillin-tazobactam, oxacillin, and linezolid (in nine patients, 16%). Moreover, 43 (81%) received steroids, 18 (34%) hydroxychloroquine, and three (6%) humanized monoclonal antibody against the interleukin-6 receptor. Routine tracheal surveillance cultures were obtained after a median of 2 days from tracheal intubation.

Among 53 tracheal aspirates, 16 (30%) were positive and 37 (70%) negative (Table 1). The causative pathogens most commonly found were *Staphylococcus aureus*, *Streptococcus pneumoniae*, and *Haemophilus influenzae*; in five cases, *Pseudomonas aeruginosa* and *Acinetobacter baumannii* were identified. Among 16 patients with a positive tracheal aspirate, nine had not received antimicrobial therapy. Empirical antimicrobial therapy was inappropriate in three of five patients with positive tracheal aspirate for pathogens not susceptible, and was titrated accordingly. Irrespective, antimicrobial therapy was continued in 21 of 37 patients (57%) with negative tracheal aspirates. Upon stratification by microbiological results of the first respiratory sample there was no difference in antimicrobial therapy, use of adjunctive anti-inflammatory therapies, and mortality.

During the 2009 H1N1 influenza pandemic, in ICU patients with a positive influenza test we routinely prescribed a neuraminidase inhibitor and empiric antibiotics against the most common causes of community-acquired pneumonia in our region (third-generation cephalosporin plus macrolide or monotherapy with respiratory fluoroquinolone). Antibiotics were rapidly de-escalated or stopped within 48 h based on the results of the first surveillance tracheal aspirate cultures.

During the early phase of the COVID-19 pandemic, when the incidence of bacterial superinfection was unknown and according to the Surviving Sepsis Campaign on the management of critically ill adults with COVID-19, empirical antibacterial therapy was administered to 49% of patients admitted in our ICU. Two recent studies reported that 70–89% of critically ill patients with COVID-19 received empiric antibiotic therapy.^{8,9} However, antimicrobial therapy was not immediately stopped after microbiological results in 57% of patients with negative tracheal surveillance aspirates with an average duration of 5 days.

The clinical confidence to limit antibiotic use after negative microbiological results was influenced by the unknown

Table 1 Clinical features of patients stratified by microbiology results of the first surveillance tracheal aspirate.*

Characteristics	All patients (53)	First positive tracheal aspirate (16)	First negative tracheal aspirate (37)	p
Male sex, % (n)	87 (46)	94 (15)	84 (31)	0.661
Age, yr	57 (51–64)	55 (53–65)	59 (47–62)	0.197
BMI, kg m ⁻²	28 (25–31)	27 (25–29)	28 (25–33)	0.313
SAPS II	33 (28–39)	31 (26–33)	34 (28–40)	0.086
Entry PEEP, cm H ₂ O	10 (10–12)	12 (10–12)	10 (10–12)	0.207
Entry P _a O ₂ /F _i O ₂ , mm Hg	113 (89–150)	109 (87–141)	115 (91–151)	0.720
Days between symptoms and ED	7 (4–10)	7 (4–9)	7 (5–10)	0.395
Days between ED and TI	3 (2–6)	5 (3–7)	3 (2–6)	0.447
Days between TI and first TA	2 (1–3)	2 (2–3)	1 (1–2)	0.053
Inpatient antimicrobial therapy, % (n pts)	48 (26)	31 (5)	57 (21)	0.135
Therapy with tocilizumab, % (n pts)	6 (3)	19 (3)	0	0.024
Therapy with OH-chloroquine, % (n pts)	34 (18)	44 (7)	30 (11)	0.358
Steroids, % (n pts)	81 (43)	94 (15)	76 (28)	0.250
C-reactive protein (mg/L)	126 (94–140)	138 (117–143)	121 (74–137)	0.132
Procalcitonin (ng/mL)	0.55 (0.18–1.21)	0.52 (0.14–0.9)	0.62 (0.38–1.8)	0.382
ICU length of stay (days)	14 (8–25)	12 (7–25)	14 (9–23)	0.635
ICU mortality, % (n)	53 (28)	62 (10)	49 (18)	0.354

ED, emergency department; SAPS II, Simplified Acute Physiology Score; TA, tracheal aspiration TI, tracheal intubation.

* Student's t-test or Mann–Whitney rank sum test, as appropriate, were used to analyse continuous variables and the χ^2 or Fisher's exact test, as appropriate, for non-continuous variables. Data are expressed as mean (standard deviation) or median (inter-quartile range) as appropriate. P<0.05.

incidence of bacterial superinfections in patients with COVID-19, by the severity of the clinical status of many patients without improvement despite the antiviral therapy, by the possibility that patients were immunocompromised because of immunomodulatory therapy, and by the fact that half of tracheal aspirates was obtained during antibiotic therapy, increasing the diagnostic difficulty.

Our preliminary data suggest that in mechanically ventilated COVID-19 patients undergoing immunomodulatory therapy, tracheal aspirates should be obtained as soon as possible and antibiotic therapy potentially withheld until microbiology results become available, because of the low rate of positive tracheal aspirates. Based on our local conditions, use of empirical broad-spectrum antibacterial drugs was inappropriate in a substantial number of cases, and narrow-spectrum antibiotics would be preferred. For use of broad-spectrum antibiotics, we can infer that de-escalation based on the results of susceptibility tests or negative tracheal aspirates should be applied as early as possible.

Absence of a specific cure, evidence that intensive care surge capabilities were rapidly overwhelmed, clinical suspicion of nosocomial infections, and wide use of immunomodulatory therapy modified our practice and led us to misuse or overuse antibacterial drugs, failing de-escalation irrespective of the culture results. Evidence from our small cohort of patients call for an urgent and comprehensive analysis of pulmonary co-infections in COVID-19 patients admitted to the ICU. In line with these goals, the COVID-19 Critical Care Consortium is currently collaborating with more than 400 ICUs worldwide to characterise secondary bacterial infections associated with SARS-CoV-2 and provide urgent recommendations on appropriate empiric therapies.

Declarations of interest

The authors declare that they have no conflicts of interest.

References

1. Huang C, Wang Y, Ren L, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506
2. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; **382**: 727–33
3. Shi H, Han X, Jiang N, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **20**: 425–34
4. Sanders JM, Monogue ML, Jodlowski TZ, Cutrell JB. Pharmacologic treatments for coronavirus disease 2019 (COVID-19): a review. *JAMA* 2020; **323**: 1824–36
5. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061–9
6. Alhazzani W, Møller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with coronavirus disease 2019 (COVID-19). *Intensive Care Med* 2020; **46**: 854–87
7. Metlay JP, Waterer GW, Long AC, et al. Diagnosis and treatment of adults with community-acquired pneumonia. An official clinical practice guideline of the American Thoracic Society and Infectious Diseases Society of America. *Am J Respir Crit Care Med* 2019; **200**: e45–67
8. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 2020; **395**: 1763–70
9. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**: 507–13

doi: 10.1016/j.bja.2020.07.019

Advance Access Publication Date: 28 July 2020

© 2020 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.

Opioids and the COVID-19 pandemic: does chronic opioid use or misuse increase clinical vulnerability?

David G. Lambert

Department of Cardiovascular Sciences (Anaesthesia), University of Leicester, Leicester, UK

E-mail: dgl3@leicester.ac.uk

Keywords: COVID-19; opioid; opioid misuse; outcome; risk factor; social determinant

Editor—Opioids have predictable analgesic actions and are widely used in many clinical settings, but they also produce unwanted side-effects including respiratory depression, tolerance and are misused. Misuse and poor opioid stewardship in the therapeutic arena are generally accepted as the underlying cause of what we describe as the ‘opioid

epidemic’ or ‘opioid crisis’.¹ According to a UK Office for National Statistics report in 2019,² there were 139 845 people in contact with drug services during the 2018–19 period; some of these will be opioid dependent, but for many this will not be a single substance misuse. Moreover, a Public Health England (PHE) report suggests 540 000 patients