



ELSEVIER

Contents lists available at ScienceDirect

Current Problems in Surgery

journal homepage: www.elsevier.com/locate/cpsurg

In Brief

Diverticulitis: An update from the age old Paradigm



Alexander T. Hawkins, MD, MPH^{a,*}, Paul E. Wise, MD^b,
 Tiffany Chan, MD, MHSc^c, Janet T. Lee, MD, MS^d,
 Tamara Glyn, MBChB^e, Verity Wood, MBChB^f,
 Timothy Eglinton, MBChB, MMedSc^e, Frank Frizelle, MBChB,
 MMedSc^e, Adil Khan, MD^g, Jason Hall, MD, MPH^h,
 M.I. Mohammed Ilyas, MDⁱ, Maria Michailidou, MD^j,
 Valentine N. Nfonsam, MD, MSⁱ, Michelle L. Cowan, MD^k,
 Jennifer Williams, MD^l, Scott R. Steele, MD, MBA^m,
 Karim Alavi, MD, MPHⁿ, C. Tyler Ellis, MD, MSCR^o,
 Danielle Collins, MD^p, Des C. Winter, MD^q, Karen Zaghiyan, MD^r,
 Gaetano Gallo, MD^s, Michele Carvello, MD^t, Antonino Spinelli, MD,
 PhD^u, Amy L. Lightner, MD^v

From the ^aSection of Colon & Rectal Surgery, Department of Surgery, Vanderbilt University Medical Center, Nashville, TN; ^bSection of Colon and Rectal Surgery, Department of Surgery, Washington University School of Medicine, St. Louis, MO; ^cUniversity of British Columbia, Vancouver, British Columbia, Canada; ^dDivision of Colon and Rectal Surgery, Department of Surgery, University of Minnesota, Saint Paul, MN; ^eDepartment of Surgery, University of Otago, Christchurch Hospital, Canterbury District Health Board, Christchurch, New Zealand; ^fChristchurch Hospital, Canterbury District Health Board, Christchurch, New Zealand; ^gRaleigh General Hospital, Beckley, WV; ^hDepartment of Surgery, Boston Medical Center, Boston, MA; ⁱDepartment of Surgery, University of Arizona, Tucson, AZ; ^jNorthern Light Health, Bangor, ME; ^kDepartment of Surgery, University of Washington, Seattle, WA; ^lEmory University, Atlanta, GA; ^mDigestive Disease and Surgery Institute, Cleveland Clinic, Cleveland, OH; ⁿDivision of Colorectal Surgery, University of Massachusetts Memorial Medical Center, Worcester, MA; ^oSection of Colon & Rectal Surgery, Hiram C. Polk, Jr., MD Department of Surgery, University of Louisville, Louisville, KY; ^pWestern General Hospital, Edinburgh, Scotland; ^qSt. Vincent's University Hospital, Dublin, Ireland; ^rCedars-Sinai Medical Center, Los Angeles, CA; ^sDepartment of Medical and Surgical Sciences, University of Catanzaro, Catanzaro, Italy; ^tColon and Rectal Surgery Unit, Humanitas Clinical and Research Center IRCCS, Department of Biomedical Sciences, Humanitas University, Milano, Italy; ^uColon and Rectal Surgery Unit, Humanitas Clinical and Research Center IRCCS, Department of Biomedical Sciences, Humanitas University, Milano, Italy; and ^vDepartment of Colorectal Surgery, Digestive Disease and Surgery Institute, Cleveland Clinic, Cleveland, OH

* Address reprint requests to Alexander T. Hawkins, MD, MPH, Section of Colon & Rectal Surgery, Vanderbilt University, 1161 21st Ave South, Room D5248 MCN, Nashville, TN 37232.

E-mail address: alex.hawkins@vumc.org (A.T. Hawkins).

<https://doi.org/10.1016/j.cpsurg.2020.100863>

0011-3840/© 2020 Elsevier Inc. All rights reserved.

For a disease process that affects so many, we continue to struggle to define optimal care for patients with diverticular disease. Part of this stems from the fact that diverticular disease requires different treatment strategies across the natural history—acute, chronic, and recurrent.

To understand where we are currently, it is worth understanding how treatment of diverticular disease has evolved. Diverticular disease was rarely described in the literature prior to the 1900s. In the late 1960s and early 1970s, Painter and Burkitt popularized the theory that diverticulosis is a disease of Western civilization based on the observation that diverticulosis was rare in rural Africa but common in economically developed countries. Previous surgical guidelines focused on early operative intervention to avoid potential complicated episodes of recurrent complicated diverticulitis (eg, with free perforation) that might necessitate emergent surgery and stoma formation. More recent data have challenged prior concerns about decreasing effectiveness of medical management with repeat episodes and the notion that the natural history of diverticulitis is progressive. It has also permitted more accurate grading of the severity of disease and permitted less invasive management options to attempt conversion of urgent operations into the elective setting, or even avoid an operation altogether.

The role of diet in preventing diverticular disease has long been debated. A high fiber diet appears to decrease the likelihood of symptomatic diverticulitis. The myth of avoid eating nuts, corn, popcorn, and seeds to prevent episodes of diverticulitis has been debunked with modern data. Overall, the recommendations for “diverticulitis diets” mirror those made for overall healthy lifestyle: high fiber, with a focus on whole grains, fruits, and vegetables.

Diverticulosis is one of the most common incidental findings on colonoscopy and the eighth most common outpatient diagnosis in the United States. More than 50% of people older than the age of 60 years and more than 60% of people older than age 80 years have colonic diverticula. Of those with diverticulosis, the lifetime risk of developing diverticulitis is estimated at 10%–25%, although more recent studies estimate a 5% rate of progression to diverticulitis. Diverticulitis accounts for an estimated 371,000 emergency department visits and 200,000 inpatient admissions per year, with an annual cost of \$2.1 to 2.6 billion per year in the United States. The estimated total medical expenditure (inpatient and outpatient) for diverticulosis and diverticulitis in 2015 was more than \$5.4 billion. The incidence of diverticulitis is increasing.

Besides increasing age, other risk factors for diverticular disease include use of nonsteroidal anti-inflammatory drugs, aspirin, steroids, opioids, smoking, and sedentary lifestyle. Diverticula most commonly occur along the mesenteric side of the antimesenteric taeniae resulting in parallel rows. These spots are thought to be relatively weak as this is the location where vasa recta penetrate the muscle to supply the mucosa. The exact mechanism that leads to diverticulitis from diverticulosis is not definitively known.

The most common presenting complaint is of left lower quadrant abdominal pain with symptoms of systemic unwellness including fever and malaise, however the presentation may vary widely. The gold standard cross-sectional imaging is multidetector computed tomography. It is minimally invasive and has sensitivity of 98% and specificity up to 99% for diagnosing acute diverticulitis. Uncomplicated acute diverticulitis may be safely managed as an outpatient in carefully selected patients. Hospitalization is usually necessary for patients with immunosuppression, intolerance to oral intake, signs of severe sepsis, lack of social support, and increased comorbidities. The role of antibiotics has been questioned in a number of randomized controlled trials and it is likely that we will see more patients with uncomplicated disease treated with observation in the future.

Acute diverticulitis can be further subclassified into complicated and uncomplicated presentations. Uncomplicated diverticulitis is characterized by inflammation limited to the colonic wall and surrounding tissue. The management of uncomplicated diverticulitis is changing. Use of antibiotics has been questioned as it appears that antibiotic use can be avoided in select groups of patients. Surgical intervention appears to improve patients' quality of life. The decision to proceed with surgery is recommended in an individualized manner.

Complicated diverticulitis is defined as diverticulitis associated with localized or generalized perforation, localized or distant abscess, fistula, stricture, or obstruction. Abscesses can be treated with percutaneous drainage if the abscess is large enough. The optimal long-term strat-

egy for patients who undergo successful nonoperative management of their diverticular abscess remains controversial. There are clearly patients who would do well with an elective colectomy and a subset who could avoid an operation altogether however the challenge is appropriate risk-stratification and patient selection. Management of patients with perforation depends greatly on the presence of feculent or purulent peritonitis, the extent of contamination, and hemodynamic status and associated comorbidities. Fistulas and strictures are almost always treated with segmental colectomy.

After an episode of acute diverticulitis, routine colonoscopy has been recommended by a number of societies to exclude the presence of colorectal cancer or presence of alternative diagnosis like ischemic colitis or inflammatory bowel disease for the clinical presentation. Endoscopic evaluation of the colon is normally delayed by approximately 6 weeks from the acute episode to reduce the risk associated with colonoscopy. Further study has questioned the need for endoscopic evaluation for every patient with acute diverticulitis. Colonoscopy should be performed routinely after complicated diverticulitis cases, when the clinical presentation is atypical or if there are any diagnostic ambiguity, or the patient has other indications for colonoscopy such as rectal bleeding or is above 50 years of age without recent colonoscopy.

For patients in whom elective colectomy is indicated, it is imperative to identify a wide range of modifiable patient comorbidities. Every attempt should be made to improve a patient's chance of successful operation. This includes optimization of patient risk factors as well as tailoring the surgical approach and perioperative management. A positive outcome depends greatly on thoughtful attention to what makes a complicated patient "complicated."

Operative management remains complex and depends on multiple factors including patient age, comorbidities, nutritional state, severity of disease, and surgeon preference and experience. Importantly, the status of surgery (ie, elective versus urgent or emergent operation) is pivotal in decisionmaking, and treatment algorithms are divergent based on the acuteness of surgery. Resection of diseased bowel to healthy proximal colon and rectal margins remains a fundamental principle of treatment although the operative approach may vary.

For acute diverticulitis, a number of surgical approaches are available, including loop colostomy, sigmoidectomy with colostomy (Hartmann's procedure), and sigmoidectomy with primary colorectal anastomosis. Overall, data suggest that primary anastomosis is preferable to a Hartman's procedure in select patients with acute diverticulitis. Patients with hemodynamic instability, immunocompromised state, feculent peritonitis, severely edematous or ischemic bowel, or significant malnutrition are poor candidates. The decision to divert after colorectal anastomosis is at the discretion of the operating surgeon. Patient factors including severity of disease, tissue quality, and comorbidities should be considered. Technical considerations for elective cases include appropriate bowel preparation, the use of a laparoscopic approach, the decision to perform a primary anastomosis, and the selected use of ureteral stents.

Management of the patient with an end colostomy after a Hartmann's procedure for acute diverticulitis can be a challenging clinical scenario. Between 20% and 50% of patients treated with sigmoid resection and an end colostomy after an initial severe bout of diverticulitis will never be reversed to their normal anatomy. The reasons for high rates of permanent colostomies are multifactorial. The debate on the best timing for a colostomy takedown continues. Six months is generally chosen as the safest time to proceed when adhesions may be at their softest, allowing for a more favorable dissection. The surgical approach will be a personal decision by the operating surgeon based on his or her experience. Colostomy takedown operations are challenging. The surgeon should anticipate and appropriately plan for a long and difficult operation. The patient should undergo a full antibiotic bowel preparation. Preoperative planning is critical; the initial operative note should be reviewed and the anatomy should be defined prior to reversal. When a complex abdominal wall closure is necessary, consideration should be given to consultation with a hernia specialist. Open surgery is the preferred surgical approach for the majority of colostomy takedown operations. Finally, ureteral catheters and diverting loop ileostomy should be considered, and the surgeon should be prepared for all anastomotic options in advance.

Since its inception in the late 1990s, laparoscopic lavage has been recognized as a novel treatment modality in the management of complicated diverticulitis and specifically for Hinchey III

(purulent) diverticulitis. Over the last decade, this treatment has been the subject of several randomized controlled trials, retrospective studies, and systematic reviews as well as cost-efficiency analyses. Despite being the subject of much debate and controversy, there is a clear role for laparoscopic lavage in the management of acute diverticulitis with the caveat that patient selection is key.

Segmental colitis associated with diverticulitis (SCAD) is an inflammatory condition affecting the colon in segments that are also affected by diverticulosis (ie, the sigmoid colon). Although SCAD is considered a separate clinical entity, it is frequently confused with diverticulitis or inflammatory bowel disease. SCAD affects approximately 1.4% of the general population and 1.15%-11.4% of those with diverticulosis and most commonly affects those in their sixth decade of life. The exact pathogenesis of SCAD is unknown, but proposed mechanisms include mucosal redundancy and prolapse occurring in diverticular segments, fecal stasis, and localized ischemia. Most cases of SCAD resolve with a high-fiber diet and antibiotics, with salicylates reserved for more severe cases. Relapse is uncommon and immunosuppression with steroids is rarely needed. A relapsing clinical course may suggest a diagnosis of inflammatory bowel disease and treatment as such should be initiated. Surgery is extremely uncommon and reserved for severe refractory disease.

Although sigmoid colon involvement is considered the most common site of colonic diverticulitis in Western countries, diverticular disease can be problematic in other areas of the colon. In Asian countries, right-sided diverticulitis outnumbers the left. This difference seems to be secondary to dietary and genetic factors. Differential diagnosis might be difficult because of similarity with appendicitis. However, accurate imaging studies allow a precise preoperative diagnosis and management planning. Transverse colonic diverticulitis is very rare, accounting for less than 1% of colonic diverticulitis with a perforation rate that has been estimated to be even more rare. Rectal diverticula are mostly asymptomatic and diagnosed incidentally in the majority of patients and rarely require treatment. Giant colonic diverticula is a rare presentation of diverticular disease of the colon and it is defined as an air-filled cystic diverticulum larger than 4 cm in diameter. The pathogenesis of giant colonic diverticula is not well defined.

Overall, the management of diverticular disease depends greatly on patient, disease, and surgeon factors. Only by tailoring treatment to the patient in front of us can we achieve optimal outcomes.