

The authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.



THE IMPORTANCE OF CORONAVIRUS DISEASE 2019 TESTING IN CARDIAC SURGERY

To the Editor:

The extraordinary demands for managing patients with Coronavirus Disease 2019 (COVID-19) has altered the Italian hospitals' ability to provide adequate care. With exponential increase of the COVID-19 population and a progressive reduction of resources, the ability to provide surgical care has been rapidly decreased.¹ Although several surgical organizations developed guides for triaging patients with cancer, the selection of candidates for cardiac surgery poses major challenges given the rapid progression of the underlying disease and the uncertain evolution of organ failure.^{2,3}

Haft and colleagues⁴ provided an important guidance to adult cardiac surgeons regarding the triage of patients with cardiac disease. The document presents some useful general rules to optimize hospital resources.⁴ However, little attention is still given to these frail patients at high risk for COVID-19 infection and dissemination. Moreover, no clear indication exists regarding the preoperative screening tests to be performed before surgery with significant variations in practice according to each institution.⁴ Our department of cardiac surgery is a tertiary referral center providing cardiac service to approximately 660,000 people in the region of Piedmont, in the northwest of Italy. Our hospital was initially COVID-19 free, but, given the large increase of infected patients, it was rapidly involved in the management of these patients. However, as per hospital policy, patients admitted for cardiac surgery were not tested for COVID-19 because they were receiving elective but not deferrable surgery. From February 1 to April 15, 100 patients were admitted to our cardiac surgery department. They were all symptomatic with a high prevalence of

New York Heart Association III or at high risk for rapidly progressive disease. At hospital admission, patients received routine laboratory tests and chest x-ray but no specific test for COVID-19. No patient showed any COVID-19-related symptoms at the time of surgery. Among operated patients, 4 of them developed symptomatic COVID-19 infection during hospitalization. One of them required intensive care unit admission. In the same time period, 5 of 8 cardiac surgeons, 1 of 9 cardiac anesthesiologists, and 7 of 48 nurses became COVID-19 positive and symptomatic. This has progressively led to a dramatic reduction in the department activity, with a 75% of reduction of the daily surgical and clinical caseload. Such dispersion of resources has consequently led to decentralization of care and a delay of surgical treatment, with 6.6% of patients who died while on the waiting list. Thus, our healthcare institutions should focus more on the triage of patients with cardiac disease, considered at high risk of COVID-19 infection. These patients should mandatorily receive microbiological tests for COVID-19, even in the presence of unstable conditions, to decrease the risk of viral dissemination and depletion of resources.

Giulia Maj, MD^a

Antonio Campanella, MD^b

Andrea Audo, MD^b

^aDepartments of Anesthesia and Intensive Care

^bCardiac Surgery

Azienda Ospedaliera SS. Antonio e Biagio e Cesare Arrigo
Alessandria, Italy

References

1. Zangrillo A, Beretta L, Silvani P, Colombo S, Scandroglio AM, Dell'Acqua A, et al. Fast reshaping of intensive care unit facilities in a large metropolitan hospital in Milan, Italy: facing the COVID-19 pandemic emergency. *Crit Care Resusc.* April 1, 2020 [Epub ahead of print].
2. COVID-19 guidance for triage of operations for thoracic malignancies: a consensus statement from thoracic surgery outcomes research network. Thoracic Surgery Outcomes Research Network, Inc. *Ann Thorac Surg.* April 4, 2020 [Epub ahead of print].
3. Drake D, Morrow CD, Kinlaw K, De Bonis M, Zangrillo A, Sade RM. Cardiothoracic surgeons in pandemics: ethical considerations. *Ann Thorac Surg.* April 9, 2020 [Epub ahead of print].
4. Haft JW, Atluri P, Alawadi G, Engelman D, Grant MC, Hassan A, et al. Adult cardiac surgery during the COVID-19 Pandemic: a tiered patient triage guidance statement. *Ann Thorac Surg.* April 10, 2020 [Epub ahead of print].

<https://doi.org/10.1016/j.jtcvs.2020.04.117>

The Editor welcomes submissions for possible publication in the Letters to the Editor section that consist of commentary on an article published in the *Journal* or other relevant issues. Authors should: • Include no more than 500 words of text, three authors, and five references. • Type with double-spacing. • See <http://jtcvs.ctsnetjournals.org/misc/ifa.html> for detailed submission instructions. • Submit the letter electronically via jtcvs.editorialmanager.com. Letters commenting on an article published in the *JTCVS* will be considered if they are received within 6 weeks of the time the article was published. Authors of the article being commented on will be given an opportunity to offer a timely response (2 weeks) to the letter. Authors of letters will be notified that the letter has been received. Unpublished letters cannot be returned.



REPLY: HAVE WE DONE THE BEST THAT WE COULD HAVE DONE?

Reply to the Editor:

In their letter to the editor,¹ Maj and colleagues cited the recently published guidance document on cardiac surgical triaging during the coronavirus 2019 (COVID-19) pandemic by Haft and colleagues² and raised the following points: (1)