

---

To claim CME credit, successfully complete this case-based posttest online at <https://digital-catalog.aad.org/diweb/catalog/t/12952/o/-esd>. Note: CME quizzes are available after the first of the month in which the article is published. If you have any questions, please contact the Member Resource Center of the American Academy of Dermatology toll-free at (866) 503-SKIN (7546), (847) 240-1280 (for international members), or by e-mailing [mrc@aad.org](mailto:mrc@aad.org).

## CME examination

Identification No. JA0820

August 2020 issue of the Journal of the American Academy of Dermatology

Creadore A, Watchmaker J, Maymone MBC, Pappas L, Lam C, Vashi NA. J Am Acad Dermatol 2020;83:315-41.

*Directions for questions 1 and 2: Choose the single best response.*

A 56-year-old white woman with history of systemic sclerosis presents to your office inquiring about treatment of facial telangiectasias. She also has microstomia and sclerodactyly.

1. Which of the following treatment options is most appropriate for her facial telangiectasias?
  - a. No treatment should be pursued given her history of scleroderma
  - b. Intense pulsed light but not pulsed dye laser
  - c. Pulsed dye laser but not intense pulsed light
  - d. 1 to 2 sessions of either intense pulsed light or pulsed dye laser
  - e. Multiple sessions of either intense pulsed light or pulsed dye laser
2. Intense pulsed light can improve what common manifestation of scleroderma?
  - a. Microstomia
  - b. Digital calcinosis
  - c. Joint contractures
  - d. Facial atrophy
  - e. Raynaud's phenomenon