

**Acral findings during the COVID-19 outbreak: Chilblain-like lesions should be preferred to acroischemic lesions**



*To the Editor:* We read with great interest the recent paper by Fernandez-Nieto et al<sup>1</sup> published in *JAAD*. The authors report a case series of acute acroischemic lesions affecting 132 nonhospitalized patients during the coronavirus disease 2019 (COVID-19) outbreak. The same cutaneous findings have been described all over the world, including a preliminary study that we conducted.<sup>2</sup>

In their article, Fernandez-Nieto et al<sup>1</sup> repeatedly use the appellative “acroischemic” for this cutaneous manifestation. Moreover, they hypothesize a relationship between a COVID-19–related altered coagulation profile and these acral lesions. As the authors state, true ischemic lesions have been reported in severely ill patients with proven coronavirus infection.<sup>3</sup>

Although the comparison between acral lesions in asymptomatic patients and ischemic lesions in severe cases is important, we find the term “acute acroischemic lesions” not accurate. Patients present with painful or itchy erythematous-edematous lesions of the extremities, sometimes evolving to blistering. This presentation is similar to what it is commonly seen in chilblains. The word “chilblains” itself etymologically refers to cold exposure (chill = cold, blain = sore). The term chilblain-like lesions, in our opinion, therefore would be preferable for the lesions that present in these patients rather than acroischemic lesions. In addition, histopathology of these lesions is quite similar to chilblains, with an absence of true necrosis; this is different to what is typically found in hospitalized patients.<sup>4</sup>

Although the exact pathogenesis of this cutaneous sign is not known yet, a worldwide common

nomenclature would in our opinion be a good starting point to avoid confusion among clinicians.

Vincenzo Piccolo, MD,<sup>a</sup> and Andrea Bassi, MD<sup>b,c</sup>

*From the Dermatology Unit, University of Campania Luigi Vanvitelli, Naples<sup>a</sup>; Unità Operativa di Dermatologia Lucca-Azienda USL Toscana Nordovest<sup>b</sup>; and the Department of Health Science, University of Florence, Florence, Italy.<sup>c</sup>*

*Funding sources:* None.

*Conflicts of interest:* None disclosed.

*IRB approval status:* Not applicable.

*Reprints not available from the author.*

*Correspondence to:* Vincenzo Piccolo, MD, c/o II Policlinico, Edificio 9, Primo piano Via Pansini 5-80131, Naples, Italy

*E-mail:* [piccolo.vincenzo@gmail.com](mailto:piccolo.vincenzo@gmail.com)

**REFERENCES**

1. Fernandez-Nieto D, Jimenez-Cauhe J, Suarez-Valle A, et al. Characterization of acute acroischemic lesions in nonhospitalized patients: a case series of 132 patients during the COVID-19 outbreak. *J Am Acad Dermatol*. 83(1):e61-e63.
2. Piccolo V, Neri I, Filippeschi C, et al. Chilblain-like lesions during COVID-19 epidemic: a preliminary study on 63 patients [e-pub ahead of print]. *J Eur Acad Dermatol Venereol*. <https://doi.org/10.1111/jdv.16526>. Accessed May 6, 2020.
3. Zhang Y, Cao W, Xiao M, et al. Clinical and coagulation characteristics in 7 patients with critical COVID-2019 pneumonia and acro-ischemia [in Chinese]. *Zhonghua Xue Ye Xue Za Zhi*. 2020;41(4):302-307.
4. Koliavas A, Dehavay F, Delplace D, et al. Coronavirus (COVID-19) infection-induced chilblains: a case report with histopathological findings. *JAAD Case Rep*. 2020;6(6):489-492.

<https://doi.org/10.1016/j.jaad.2020.05.077>