FISEVIER


Contents lists available at ScienceDirect

Annals of Diagnostic Pathology

journal homepage: www.elsevier.com/locate/anndiagpath



Retraction notice





Koichi Sato^a, Takeshi Uehara^b, Mai Iwaya^b, Tomoyuki Nakajima^b, Yusuke Miyagawa^c, Tomoaki Suga^a, Hiroyoshi Ota^d, Eiji Tanaka^a

- ^a Department of Gastroenterology, Shinshu University School of Medicine, Matsumoto, Japan
- b Department of Laboratory Medicine, Shinshu University School of Medicine, Matsumoto, Japan
- ^c First Department of Surgery, Shinshu University School of Medicine, Matsumoto, Japan

This article has been retracted: please see Elsevier Policy on Article Withdrawal (https://www.elsevier.com/about/our-business/policies/article-withdrawal).

This article has been retracted at the request of the authors regret an error in the calculation with the statistical software (JMP ver.12). The survival status of overall survival was set as dead = 1, alive = 0 when

performing log-rank test and univariate analysis with statistical software, but the initial setting in JMP is dead =0, alive =1, so the incorrect result was published. After the correction, there is no statistical significance between LGR5 expression and the Log-rank test and the univariate analysis. The authors would like to apologize for any inconvenience caused.

d Department of Biomedical Laboratory Medicine, Shinshu University School of Medicine, Matsumoto, Japan