

Associate Editorial Board

Tom Abbott, Royal London Hospital, London, UK

Mohamed Abdulatif, Cairo University, Cairo, Egypt

Michael Aziz, Oregon Health & Science University, Portland, OR, USA

Tim Cook, Royal United Hospital, Bath, UK

Andrea Cortegiani, University of Palermo, Palermo, Italy

Melissa Cushing, Weill Cornell Medicine, New York, NY, USA

Judith Dinsmore, St George's Hospital, London, UK

Talmage Egan, University of Utah School of Medicine, Salt Lake City, UT, USA

Matthias Eikermann, Harvard Medical School, Boston, MA, USA

Thomas Engelhardt, Montreal Children's Hospital, Montreal, Canada

Lis Evered, St Vincent's Hospital Melbourne, Australia

Dominique Fletcher, Raymond Poincaré Hospital Garches, Paris, France

Patrice Forget, University of Aberdeen, Aberdeen, UK

Ruth Gotian, Weill Cornell Medical College, New York, NY, USA

Charles Hogue, Northwestern University School of Medicine, Chicago, IL, USA

Vesna Jevtovic-Todorovic, University of Colorado, Denver, CO, USA

Keyvan Karkouti, University of Toronto, Ontario, Canada

Massimo Lamperti, Cleveland Clinic Abu Dhabi, UAE

Andrew Lumb, Leeds Teaching Hospitals, Leeds, UK

Alan Macfarlane, Glasgow Royal Infirmary, Glasgow, UK

Keira Mason, Harvard Medical School, Boston Children's Hospital, Boston, MA, USA

Colin McCartney, University of Toronto, Ontario, Canada

Graeme McLeod, Ninewells Hospital and Medical School, Dundee, UK

Alexander Ng, Royal Wolverhampton Hospitals NHS Trust, Wolverhampton, UK

Barbara J. Philips, St George's, University of London, London, UK

Kane Pryor, Weill Cornell Medical College, New York, NY, USA

Ben Shelley, University of Glasgow, Glasgow, UK

Jamie Sleigh, University of Auckland, Hamilton, New Zealand

Ken Tanaka, University of Maryland, Baltimore, MD, USA

Vivianne L. Tawfik, Stanford University School of Medicine, Stanford, CA, USA

Laszlo Vutskits, University Hospitals of Geneva, Geneva, Switzerland

Joyce Wahr, University of Minnesota, Minneapolis, MN, USA

Vishvarani Wanigasekera, University of Oxford, Oxford, UK

Christopher Wu, Hospital for Special Surgery, New York, NY, USA

Advisers on statistics:

Malachy Columb, Wythenshawe Hospital, Manchester, UK

David Zurakowski, Harvard Medical School, Boston Children's Hospital, Boston, MA, USA

Cover image

Perioperative fluid management has an important impact on patient outcomes such that optimal fluid management should reduce complications. By analysing a large US database, Miller and colleagues (pp 720–29) conclude that either very high or very low perioperative fluid volumes is associated with more complications after noncardiac surgery.

Credit: 123rf.com; copyright: Thirasak Phuchom