

follow-up. COVID-19 ICU survivors will need continued follow-up and support to characterise these psychological sequelae and to help mitigate the effects of this significant life event.

'Long Covid' has been used to describe symptoms in people reporting long-term effects after COVID-19, but we would urge caution in applying this diagnosis to ICU survivors who may simply be experiencing the symptoms and recovery typical of many ARDS survivors.

We have observed significant physical weakness in critically ill patients recovering from COVID-19, highlighting the need for ongoing physical rehabilitation in this patient group. Detailed analysis of both ICU care and follow-up of COVID-19 patients may allow identification of the most favourable management strategies of patients with severe COVID-19 in order to mitigate long-term sequelae.

A limitation of this study is that it is a single-centre review; therefore, our findings may not reflect the outcomes of patients cared for in other ICUs. Furthermore, the number of survivors in whom we report data is relatively small. However, our admission characteristics and detailed descriptors of ICU stay allow other units to make a comparison with their own data. Our results may assist in health service planning and ongoing care and support requirements for an ever-increasing number of mechanically ventilated COVID-19 survivors.

Declarations of interest

GK is an editorial fellow of *British Journal of Anaesthesia*. No other conflicts of interest exist.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bja.2021.03.005>.

References

1. ICNARC – ICNARC report on COVID-19 in critical care: England, Wales and Northern Ireland 19 February 2021. Available from: <https://www.icnarc.org/Our-Audit/Audits/Cmp/Reports> (accessed 24 Feb 2021).
2. van den Borst B, Peters JB, Brink M, et al. Comprehensive health assessment three months after recovery from acute COVID-19. *Clin Infect Dis* 2020; ciaa1750. <https://doi.org/10.1093/cid/ciaa1750>. Advanced access published on November 21. [update]
3. Arnold DT, Hamilton FW, Milne A, et al. Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: results from a prospective UK cohort. *Thorax* 2020. <https://doi.org/10.1136/thoraxjnl-2020-216086>. Advanced access published on December 03. [update]
4. Taboada M, Moreno E, Cariñena A, et al. Quality of life, functional status, and persistent symptoms after intensive care of COVID-19 patients. *Br J Anaesth* 2021; 126: e110–3
5. Tabatabaei SMH, Rajebi H, Moghaddas F, Ghasemiadl M, Talari H. Chest CT in COVID-19 pneumonia: what are the findings in mid-term follow-up? *Emerg Radiol* 2020; 27: 711–9
6. Massy-Westropp NM, Gill TK, Taylor AW, Bohannon RW, Hill CL. Hand grip strength: age and gender stratified normative data in a population-based study. *BMC Res Notes* 2011; 4: 127
7. Chiumello D, Coppola S, Froio S, Gotti M. What's next after ARDS: long-term outcomes. *Respir Care* 2016; 61: 689–99
8. Guler SA, Ebner L, Beigelman C, et al. Pulmonary function and radiological features four months after COVID-19: first results from the national prospective observational Swiss COVID-19 lung study. *Eur Respir J* 2021. <https://doi.org/10.1183/13993003.03690-2020>. Advanced access published on January 8. [update]
9. Dowdy DW, Eid MP, Dennison CR, et al. Quality of life after acute respiratory distress syndrome: a meta-analysis. *Intensive Care Med* 2006; 32: 1115–24

doi: 10.1016/j.bja.2021.03.005

Advance Access Publication Date: 19 March 2021

Crown Copyright © 2021 Published by Elsevier Ltd on behalf of British Journal of Anaesthesia. All rights reserved.

Rationing oxygen use during total intravenous anaesthesia: a proportionate response?

Morgan Back^{1,*}, George Zhong², Adam Al-Attar¹, Rebecca Sutton¹ and Cliff Shelton^{1,3}

¹Department of Anaesthesia, Manchester University NHS Foundation Trust, Manchester, UK, ²Department of Anaesthesia, Concord Repatriation General Hospital, Sydney, Australia and ³Lancaster Medical School, Faculty of Health and Medicine, Lancaster University, Lancaster, UK

*Corresponding author. E-mail: morganback@gmail.com

Keywords: COVID-19; fresh gas flow; healthcare cost; sustainability; total intravenous anaesthesia

Editor—We read with interest Hall and Chakladar's¹ comment on the recent study by Zhong and colleagues²

of the environmental and economic impacts of fresh gas flow (FGF) during total i.v. anaesthesia. Their concern

about oxygen use during the COVID-19 pandemic sadly remains topical, as the disease continues to overwhelm critical care capacity in many hospitals.

Hall and Chakladar¹ suggest that higher FGF rates during the provision of non-inhalation general anaesthesia should be avoided until after the pandemic has abated, citing reports of hospitals that have experienced oxygen shortages.¹ In the UK, which continues to be severely affected by COVID-19, there have been several cases of oxygen supply issues during surges of hospital admissions. However, this appears not to be as a result of a lack of oxygen *per se*, but the per minute oxygen demand exceeding the flow capacity of supply systems.³ Internationally, absolute shortages of oxygen have been experienced, for example in sub-Saharan Africa.⁴ This appears to be largely as a result of economic and infrastructure factors, but is not representative of a global oxygen shortage.

Whilst low-flow anaesthesia should form part of a strategy to cope with exceptional surges in oxygen demand, its potential contribution to this endeavour is unfortunately minimal. The total consumption of oxygen per patient using higher FGFs via a circle system is not as great as it may at first appear when considered in context. Using Hall and Chakladar's¹ example, a 6-h case using FGF of 6 L min⁻¹ and a fraction of inspired oxygen (FiO₂) of 0.3 requires 252 L of oxygen. Whilst this may appear to be a large volume, it represents only 0.7 L min⁻¹ of oxygen gas usage; lower than the minimal requirements of even low-flow nasal cannulae.

Furthermore, it has been shown that FiO₂ may be significantly lower than the fraction of oxygen delivered at low FGFs because of both oxygen consumption and increased rebreathing of exhaled gases by the patient.⁵ While Hall and Chakladar¹ suggest that providing an FiO₂ of 0.3 at 1 L min⁻¹ requires an oxygen flow of 0.1 L min⁻¹, Hendrickx and colleagues⁶ found in their *in vivo* study that oxygen flows of 0.2 L min⁻¹ were required to maintain a steady-state FiO₂ of 0.31 within the circle system at a total FGF of 1 L min⁻¹. It is likely that even higher oxygen flow rates would be required for patients with increased oxygen consumption at low FGFs. Consequently, oxygen savings are not proportional to the reduction in FGF for a given FiO₂.

During times of extraordinary oxygen demand we agree that clinicians may be required to take all available steps to safely minimise consumption. But even during the COVID-19 pandemic, increasing the FGF from 1 to 6 L min⁻¹ during non-inhalation anaesthesia remains a reasonable option at most times, representing only a 0.5 L min⁻¹ increase in oxygen gas usage when delivering an FiO₂ of 0.3.

In addition to its tragic impacts on the health and wellbeing of millions of people, the COVID-19 pandemic is responsible for profound adverse economic and environmental effects.^{7,8} Measures that clinicians can take to mitigate these problems, including optimising FGF rates, are perhaps now more relevant than ever.⁹

Declarations of interest

CS is a former member of the editorial board of *BJA Education*. The other authors declare that they have no conflicts of interest.

References

- Hall A, Chakladar A. High fresh gas flow during non-inhalational anaesthesia during the COVID-19 pandemic. *Comment on Br J Anaesth* 2020; 125: 773-778. *Br J Anaesth* 2020; 126: e38-9
- Zhong G, Abbas A, Jones J, Kong S, McCulloch T. Environmental and economic impact of using increased fresh gas flow to reduce carbon dioxide absorbent consumption in the absence of inhalational anaesthetics. *Br J Anaesth* 2020; 125: 773-8
- Collision H. UK's Hancock: National oxygen supplies aren't running short. Available from: <https://www.politico.eu/article/uks-hancock-national-oxygen-supplies-arent-running-short/> (accessed 9 February 2021).
- Stein F, Perry M, Banda G, Woolhouse M, Mutapi F. Oxygen provision to fight COVID-19 in sub-Saharan Africa. *BMJ Glob Health* 2020; 5: e002786
- Herbert L, Magee P. Low flow anaesthesia and circle systems. *BJA Educ* 2017; 17: 301-5
- Hendrickx J, De Cooman S, Vandepuit D, et al. Air-oxygen mixtures in circle systems. *J Clin Anesth* 2001; 13: 461-4
- White SM, Shelton CL. Effects of the COVID-19 pandemic on environmental sustainability in anaesthesia. *Br J Anaesth* 2021; 126: e118-9
- McGain F, Muret J, Lawson C, Sherman JD. Effects of the COVID-19 pandemic on environmental sustainability in anaesthesia. *Response to Br J Anaesth* 2021; 126: e118-e119. *Br J Anaesth* 2021; 126: e119-22
- Back M, Al-Attar A, Sutton R, Shelton CL. Fresh gas flow during total intravenous anaesthesia and marginal gains in sustainable healthcare. *Comment on Br J Anaesth* 2020; 125: 773-8. *Br J Anaesth* 2021; 126: e143-4

doi: 10.1016/j.bja.2021.02.031

Advance Access Publication Date: 19 March 2021

© 2021 Published by Elsevier Ltd on behalf of British Journal of Anaesthesia.

Protection of healthcare workers during aerosol-generating procedures with local exhaust ventilation

Jante S. Sinnige, Fabian O. Kooij, Hans van Schuppen, Markus W. Hollmann* and Nicolaas H. Sperna Weiland

Department of Anaesthesiology, Amsterdam UMC, Amsterdam, the Netherlands

*Corresponding author. E-mail: m.w.hollmann@amsterdamumc.nl