

References

1. Kaushik S, Aydin SI, Derespina KR, Bansal PB, Kowalsky S, Trachtman R, et al. Multisystem inflammatory syndrome in children associated with severe acute respiratory syndrome coronavirus 2 infection (MIS-C): a multi-institutional study from New York City. *J Pediatr* 2020;224:24-9.
2. Carter MJ, Fish M, Jennings A, Doores KJ, Wellman P, Seow J, et al. Peripheral immunophenotypes in children with multisystem inflammatory syndrome associated with SARS-CoV-2 infection. *Nat Med* 2020;26:1701-7.
3. Goldstein MR, Poland GA, Graeber CW. Does apolipoprotein E genotype predict COVID-19 severity? *QJM* 2020;113:537-8.
4. Kuo CL, Pilling LC, Atkins JL, Masoli JAH, Delgado J, Kuchel GA, et al. ApoE E4 genotype predicts severe COVID-19 in the UK Biobank Community Cohort. *J Gerontol A Biol Sci Med Sci* 2020;20:1-2.
5. Mahley RW, Weisgraber KH, Huang Y. Apolipoprotein E: structure determines function, from atherosclerosis to Alzheimer's disease to AIDS. *J Lipid Res* 2009;50:S183-8.
6. Gale SC, Gao L, Mikacenic C, Coyle SM, Rafaels N, Dudenkov TM, et al. *APOE4* is associated with enhanced in vivo innate immune responses in human subjects. *J Allergy Clin Immunol* 2014;134:127-34.
7. Corbo RM, Scacchi R. Apolipoprotein E (APOE) allele distribution in the world. Is *APOE*4* a 'thrifty' allele? *Ann Hum Genet* 1999;63:301-10.

RNA is not virus



To the Editor:

Yonker et al demonstrated high levels of viral RNA in nasopharyngeal samples obtained from children.¹ A plausible explanation for the seemingly counterintuitive lack of symptoms and high viral load is the methods employed are detecting high viral RNA, which does not correlate to high viral load as the authors suggest. Establishing a correlation between RNA and virus in the asymptomatic pediatric population with a virus culture is a prerequisite for assertion that RNA positive children carry high viral loads.² An alternative explanation is that the children in the study, and generally most children, mount an effective immune response to subclinical infection. That immune response causes lysis of infected cells, spilling cellular contents including viral RNA and proteins into surrounding tissue or interference with viral assembly inhibiting excretion of infectious virus. High RNA in nasopharyngeal samples obtained from asymptomatic patients is a reflection of an effective immune response rather than high viral load, explaining minimal infectivity of such subjects.³

Gregory I. Sawchyn, MD, MBA

Population Health
Sound Physicians Group
Tacoma, Washington

<https://doi.org/10.1016/j.jpeds.2020.09.057>

The author declares no conflicts of interest.

References

1. Yonker LM, Neilan AM, Bartsch Y, Patel AB, Regan J, Arya P, et al. Pediatric severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2): clinical presentation, infectivity, and immune responses. *J Pediatr* 2020;227:45-52.e5.
2. La Scola B, Le Bideau M, Andreani J, Hoang VT, Grimaldier C, Colson P, et al. Viral RNA load as determined by cell culture as a management tool for discharge of SARS-CoV-2 patients from infectious disease wards. *Eur J Clin Microbiol Infect Dis* 2020;39:1059-61.
3. Kim J, Choe YJ, Lee J, Park YJ, Park O, Han MS, et al. Role of children in household transmission of COVID-19. *Arch Dis Child*, in press.

Reply



To the Editor:

In our article, we report high viral load in children infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the first 2 days of illness. We did not assess the viability of virus from these respiratory secretions and agree that RNA does not necessarily correlate with live virus. However, in hospitalized adults, live virus is readily cultured from respiratory secretions during the first week of symptoms.¹ It is plausible that differences exists between pediatric and adult immune responses to SARS-CoV-2²; distinctions in mucosal immune responses could impact viral detection by reverse-transcription polymerase chain reaction and/or severity of symptoms. Another plausible hypothesis is that SARS-CoV-2 could colonize the upper airways efficiently in both children and adults but children may be less likely to have colonization in the lower respiratory tract. Although we described SARS-CoV-2 serology and show that infected children mount a humoral response following infection, we did not assess mucosal responses or regional differences of viral load within the airways. Research is needed to test these hypotheses.

Lael M. Yonker, MD

Alessio Fasano, MD

Mucosal Immunology and Biology Research Center
Department of Pediatrics
Massachusetts General Hospital
Harvard Medical School
Boston, Massachusetts

<https://doi.org/10.1016/j.jpeds.2020.09.058>

References

1. Wolfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Muller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020;581:465-9.
2. Fialkowski A, Gernez Y, Arya P, Weinacht KG, Kinane TB, Yonker LM, et al. Insight into the pediatric and adult dichotomy of COVID-19: age-related differences in the immune response to SARS-CoV-2 infection. *Pediatr Pulmonol* 2020;55:2556-64.